

# **2024 WILD TURKEY SUMMER OBSERVATION SURVEY REPORT**

Hannah Plumpton  
Upland Game Bird Biologist

## **Survey Overview**


Each summer, the North Carolina Wildlife Resources Commission (NCWRC) coordinates an observation survey to gain insight into wild turkey productivity and carryover of gobblers from the previous spring turkey hunting season. This year survey cards were mailed to approximately 5,000 people. The mailing list included a mix of NCWRC employees, National Wild Turkey Federation members, and other individuals that had participated in the survey previously. Several news releases and social media posts were used to recruit new participants this year. An unknown number of new participants were able to access the online survey using their existing WRC ID number. Additionally, our support staff helped several new participants establish their WRC ID number to begin reporting sightings of turkeys.

As in previous years, participants reported wild turkeys they observed during the course of routine daily activities from July 1<sup>st</sup> through August 31<sup>st</sup>. Participants recorded observations in all of North Carolina's 100 counties. There were 18 counties that had fewer than 10 participants and 16 counties that had fewer than 25 observations (Figures 1 and 2). A total of 1,244 individuals participated in the survey in 2024. They recorded a total of 7,413 separate observations and saw a combined total of 38,363 wild turkeys (Table 1). This is an increase from last year's survey when 1,276 participants reported 7,264 observations. Participants this year reported 6,782 observations via the traditional survey cards and 631 observations via the on-line application. At current participation levels, the summer observation provides meaningful insight into our wild turkey population and offers a way to gauge hunting pressure and population trends across the state.


## **Data Analysis**

As in previous years, the data were compiled, checked for errors, and analyzed to determine a productivity index from poult per hen ratios and to evaluate carryover of gobblers from gobblers per hen ratios. Estimates of productivity were derived from the ratios of poults and hens in each reported observation, rather than from the total number of hens and poults observed. This approach recognizes the fact that the reported turkey observations are just a sample of the entire population and that a measurement of error is part of the estimation process. Specifically, this approach provides a way to compute a 95% confidence interval for each estimate. The large number of participants and observations in this survey allows for precise estimates at regional scales, hence the relatively small

confidence intervals in Table 2 and Figures 3, 4, and 5. Gobblers per hen ratios were calculated based on the sum of all observations.



**Figure 1.** Number of participants reporting turkeys in each county in the 2024 Wild Turkey Summer Observation Survey. Some participants reported turkeys from more than one county.



**Figure 2.** Number of observations reported in each county in the 2024 Wild Turkey Summer Observation Survey.

<b>Region*</b>	<b>Observations</b>	<b>Hens w/o Poults</b>	<b>Hens w/ Poults</b>	<b>Total Hens</b>	<b>Total Poults</b>	<b>Total Gobblers</b>	<b>Total Unk.</b>
<b>Coastal</b>	2,875	2,942	1,901	4,843	5,247	2,654	3,537
<b>Piedmont</b>	2,399	2,161	1,360	3,521	4,171	1,611	1,964
<b>Mountain</b>	2,139	2,144	1,382	3,526	4,224	1,365	1,700
<b>State</b>	<b>7,413</b>	<b>7,247</b>	<b>4,643</b>	<b>11,890</b>	<b>13,642</b>	<b>5,630</b>	<b>7,201</b>

\*Geographical regions, not NCWRC regions.

## **Productivity**


Wild turkey productivity can be evaluated by examining the observations of hens and poults in the survey. This information is best considered in a relative fashion, comparing the data among the three regions and also evaluating the trends through time. There are three primary ways to evaluate productivity:

- Poults per Hen – this ratio gives an indication of overall productivity
- Poults per Brood – this ratio gives an indication of poult survival
- Percentage of Hens Observed with Poults – indicates nesting success


Productivity statewide was estimated to be 1.5 poults per hen and was higher in the mountain and piedmont regions than the coastal region (Table 2). Productivity was 1.6 poults per hen in the mountain region and 1.5 and 1.4 poults per hen in the piedmont and coast, respectively. From a biological standpoint, these estimates are comparable. Productivity in 2024 was greater than in 2023 and slightly lower than in 2022 (Figure 3). Poult survival statewide was 3.4 poults per brood and was noticeably lower in the coastal region compared to the piedmont and mountain region. The percentage of hens observed with poults was consistent across the state. Our estimates of turkey reproduction this year are on par with what we have observed over the course of the last decade (Figures 4 and 5).

<b>Region*</b>	<b>% Hens with Poults</b>	<b>Poults Per Brood</b>	<b>Poults Per Hen</b>	<b>Gobblers/Hen Ratio</b>
<b>Coastal</b>	39%	3.2 (3.1 – 3.3)	1.4 (1.3 -1.5)	0.55
<b>Piedmont</b>	39%	3.5 (3.4 – 3.6)	1.5 (1.4 – 1.6)	0.46
<b>Mountain</b>	39%	3.5 (3.4 – 3.7)	1.6 (1.5 – 1.7)	0.39
<b>State</b>	<b>39%</b>	<b>3.4 (3.3 – 3.5)</b>	<b>1.5 (1.4 – 1.6)</b>	<b>0.47</b>


\*Geographical regions, not NCWRC regions.



**Figure 3.** Regional productivity estimates from 2022–2024 Wild Turkey Summer Observation Survey. Error bars represent 95% confidence intervals.



**Figure 4.** Statewide productivity estimates from Wild Turkey Summer Observation Surveys, 2015–2024. Error bars represent 95% confidence intervals.




**Figure 5.** Statewide poult survival estimates from Wild Turkey Summer Observation Surveys, 2015-2024. Error bars represent 95% confidence intervals.

### **Gobbler Carryover**

The observed ratio of gobblers per hen indicates the level of carryover of gobblers from the previous spring turkey hunting season. Higher levels of gobbler harvest by hunters will typically result in lower gobblers per hen ratios. A ratio of less than 0.50 gobblers per hen may be an indication of over-harvest of the male segment of the turkey population if quality spring gobbler hunting is the management goal.

Over the past 10 years, gobblers per hen ratios in the summer observation survey have been between 0.47 and 0.57 gobblers per hen (Figure 6). The ratio for the 2024 summer observation survey was 0.47 gobblers per hen. These data indicate that, if quality spring gobbler hunting is to be maintained, additional pressure should not be placed on the male segment of the wild turkey population by increasing the season length, opening the spring season earlier, or increasing the bag limit.



**Figure 6.** Ratio of gobblers per hen observed in Wild Turkey Summer Observation Surveys, 2015-2024.